

A generalization of the Shapley-Ichiishi result

Jeroen Kuipers ^{*} Dries Vermeulen [†] Mark Voorneveld ^{‡§}

March 3, 2009

Abstract

The Shapley-Ichiishi result states that a game is convex if and only if the convex hull of marginal vectors equals the core. In this paper we generalize this result by distinguishing equivalence classes of balanced games that share the same core structure. We then associate a system of linear inequalities with each equivalence class, and we show that the system defines the class. Application of this general theorem to the class of convex games yields an alternative proof of the Shapley-Ichiishi result. Other applications range from computation of stable sets in non-cooperative game theory to determination of classes of TU games on which the core correspondence is additive (even linear). For the case of convex games we prove that the theorem provides the minimal defining system of linear inequalities. An example shows that this is not necessarily true for other equivalence classes of balanced games.

SSE/EFI Working Paper Series in Economics and Finance, No. 711.

^{*}MICC-Mathematics, Maastricht University, P.O.Box 616, 6200 MD Maastricht, The Netherlands.

[†]Department of Quantitative Economics, Maastricht University, P.O.Box 616, 6200 MD Maastricht, The Netherlands.

[‡]Department of Economics, Stockholm School of Economics, P.O.Box 6501, 113 83 Stockholm, Sweden. Financial support from the Wallander/Hedelius Foundation is gratefully acknowledged.

[§]We thank two anonymous referees and William Thomson for their helpful comments on earlier drafts of this paper and Jean Derks for many fruitful discussions.

1 Introduction

In his study of convex games, Shapley (1971) proved that the core of a convex game is the convex hull of the marginal vectors of that game. A decade later, Ichiishi (1981) showed that the converse of this statement is also true: If the core of a game is equal to the convex hull of marginal vectors, then the game is convex. The combined Shapley-Ichiishi result thus relates the defining system of linear inequalities of convex games to the core structure that all convex games share. The interesting theoretical properties of convex games has led to a vast literature in which convex games play a role. Examples are unanimity games, bankruptcy games (O'Neill, 1982, Aumann and Maschler, 1985), sequencing games (Curiel *et al.*, 1989), standard tree games (Megiddo, 1978, Granot *et al.*, 1994), and extended tree games (Granot *et al.*, 2002). The Shapley-Ichiishi result is used freely in the analysis of such games.

In this paper, we generalize the Shapley-Ichiishi result to other classes of balanced games. Specifically, we distinguish equivalence classes of balanced games with the same core structure, and we provide a procedure which, given such an equivalence class, generates a defining system of linear inequalities that describes the closure of the class. One of the equivalence classes is the class of strictly convex games, and its closure is the class of convex games.

The main motivation for this paper comes from the literature on strategic stability for bimatrix games. Hillas (1990) defined quasi-stable sets. A closely related notion, in fact quasi-stability minus the invariance and minimality requirements, is the notion of a Q-set, defined in Vermeulen *et al.* (1996). For a bimatrix game (A, B) , a Q-perturbed version of (A, B) is a strategic form game in which the payoffs for the players are the same as in the original game, but each player is only allowed to select strategies from a strict-subset of his strategy space. These subsets take the form of the core of a TU-game, which gives the connection to the current paper. A closed set S of strategy pairs is a Q-set if every Q-perturbed version of (A, B) has an equilibrium close to S . Vermeulen *et al.* (1996) showed that for bimatrix games, minimal Q-sets are finite.

Their approach also shows that, given a closed set S of strategy pairs for a given bimatrix game, testing whether S is a Q-set is a finite task. The reason for this is that many Q-perturbed versions of (A, B) in fact impose the same conditions on S for it to be a Q-set. That way we can construct an equivalence relation on Q-perturbed versions of (A, B) , by saying that two Q-perturbations are equivalent when they impose the same condition

on S . It is shown in Vermeulen *et al.* (1996) that there are only finitely many equivalence classes under this equivalence relation.

Now, the strategy spaces of two equivalent Q -perturbations, viewed as cores of two TU-games, can be shown to have the same core structure. Thus, when devising a finite test to check whether a closed set S is a Q -set, one needs to determine all classes of TU-games having the same core structure. The search for classes of TU-games having the same core structure becomes manageable when these classes are given by linear (in)equalities. And the task to describe all classes of TU-games having the same core structure by means of linear (in)equalities is executed in this paper. In that sense the current paper develops one of the tools needed to construct a finite test for Q -sets.

A second application is analysis of the core correspondence. As is well known, the core correspondence is superadditive. The equivalence classes defined in this paper are exactly those sets of TU games on which the core is precisely additive. These sets can be determined by computing the linear inequalities that are presented in our main theorem.

A third application of the generalized Shapley-Ichiishi result is the analysis of certain classes of games defined by linear inequalities, for example the class of k -convex games. It is known that these games are balanced and that a full characterization of the extreme core elements can be given (Driessen (1986)). Indeed, this characterization implies that all k -convex games share the same core structure, as we define it in Section 2 of this paper. Therefore, the main theorem of our paper also applies to the class of k -convex games.

The setup of this paper is as follows. In Section 2, we define the core structure of a game, which gives rise to equivalence classes of games with the same core structure, and we describe the core structure of convex games. In Section 3, we give a short review on polyhedra and their facial structure. In Section 4, we provide a constructive method for generating a system of linear inequalities associated with a given core structure, and we show that the system defines the closure of the equivalence class. An example shows that the system may contain redundant inequalities. From the description with inequalities, it follows that the closure is a polyhedral cone. These cones are of interest from a computational perspective, as it can be shown that the core is an additive correspondence on each cone. In Section 5, we apply our main theorem to the class of convex games, and this way, we derive an alternative proof of the Shapley-Ichiishi result. In fact, we obtain a somewhat sharper result, since the set of defining inequalities that

we obtain turns out to be a strict subset of the set of inequalities that is normally used to define convexity. We then prove that each inequality in our system describes a facet of the cone of convex games. Hence, in the case of convex games, we obtain the minimal system.

2 Games and their core structure

A transferable utility game, or *game*, is a pair (N, v) where $N = \{1, \dots, n\}$ is the set of *players* and v is a function that assigns to each coalition $S \subseteq N$ its *worth* $v(S) \in \mathbb{R}$. The worth $v(\emptyset)$ of the empty set is zero. Throughout this paper we keep the player set N fixed. So we simplify notation and write v instead of (N, v) to denote a game.

A vector $x \in \mathbb{R}^N$ is an *allocation*. The i^{th} coordinate x_i of the allocation x represents the payoff to player $i \in N$. For coalition $S \subseteq N$, the aggregate payoff $\sum_{i \in S} x_i$ is denoted by $x(S)$. An allocation x is *efficient for v* if it distributes the worth of the grand coalition among the players of the game v , i.e., if $x(N) = v(N)$. An efficient allocation x is a *core allocation for v* if

$$x(S) \geq v(S) \text{ for all } S \subset N.$$

The set of core allocations, denoted by $C(v)$, is called the *core* of v . A game whose core is not empty is *balanced*.

A nonempty coalition S with $x(S) = v(S)$ is called *tight at x in v* . The collection of coalitions that are tight at x in v is denoted by $T(v, x)$. Define the *core structure of v* by

$$\mathcal{T}(v) := \{T(v, x) \mid x \text{ is an extreme point of } C(v)\}.$$

We say that w is a *limit game for v* if for every extreme point $x \in C(v)$ there exists an extreme point $y \in C(w)$ such that $T(v, x) \subseteq T(w, y)$. The collection of limit games for v is denoted by $L(v)$.

In the remaining part of this section, we focus on convex games and their core structure. A game v is *convex* if it satisfies the inequalities

$$v(S) + v(T) \leq v(S \cap T) + v(S \cup T) \tag{*}$$

for all coalitions S and T . If the inequality in $(*)$ is strict whenever $S \not\subseteq T$ and $T \not\subseteq S$, the game is *strictly convex*.

Ichiishi (1981) showed that a game v is convex if and only if it satisfies the increasing marginal contributions (imc) inequalities

$$v(T \cup \{i\}) - v(T) \geq v(S \cup \{i\}) - v(S) \quad (**)$$

for all $S \subseteq T$ and $i \notin T$. So, not all inequalities in (*) are needed, and (**) suffices. However, also the system (**) has redundant inequalities, as will become apparent later on in this paper.

One of the main results on convex games is the Shapley-Ichiishi theorem. It can be stated as follows. Let v be a game. Consider for a given permutation σ of the player set N the corresponding marginal vector $m^\sigma(v)$ defined by

$$m^\sigma(v)_{\sigma(k)} := v(S_k^\sigma) - v(S_{k-1}^\sigma)$$

for each player $k \in N$, where $S_0^\sigma := \emptyset$ and

$$S_k^\sigma := \{\sigma(j) \mid j \leq k\}$$

for each $k \in \{1, \dots, n\}$. The result of Shapley (1971) and Ichiishi (1981) is

Theorem 1 *The game v is convex if and only if $m^\sigma(v) \in C(v)$ for each permutation σ .*

For a convex game v , Theorem 1 implies that $\{S_k^\sigma \mid k \in N\} \subseteq T(v, m^\sigma)$ for each permutation σ . For a strictly convex game v , Shapley (1971) proved that $T(v, m^\sigma) = \{S_k^\sigma \mid k \in N\}$ for each permutation σ . Hence, the core structure of a strictly convex game is given by

$$\{\{S_k^\sigma \mid k \in N\} \mid \sigma \text{ is a permutation of } N\},$$

and every convex game is a limit game for a given strictly convex game.

3 The facial structure of polyhedra

Let A be an $m \times n$ matrix and let $b \in \mathbb{R}^m$. The solution set

$$P = \{x \in \mathbb{R}^n \mid Ax \geq b\}$$

of the system of linear inequalities $Ax \geq b$ is a *polyhedron*. If $b = 0$, then obviously P is a cone, and P is called a *polyhedral cone*. Evidently, every polyhedron is a convex set. The system $Ax \geq b$ is a *defining system for P* .

The dimension of a convex subset C of \mathbb{R}^n is defined as the dimension of the smallest affine subspace that contains C . It is denoted by $\dim(C)$. A closed subset F of C is a *face* of C if $x, y \in F$ if and only if $\frac{1}{2}x + \frac{1}{2}y \in F$. This definition implies that a face is convex. Thus each face has a dimension and a face with dimension $\dim(C) - 1$ is called a *facet*. A zero-dimensional face is called an *extreme point*.

For polyhedra, there is an alternative way to describe faces. Denote the i -th row of A by a_i , and let I denote an index set of rows of A . Define

$$F_I = \{x \in P \mid a_i x = b_i \text{ for all } i \in I\}.$$

It is easy to prove that each F_I is a face of P , and conversely (see e.g. Schrijver (1986)) each face of P is of the type F_I . The polyhedron P itself is obtained when we adopt the convention $P_\emptyset = P$. Every extreme point of P can be obtained from an index set I , such that the rows a_i for $i \in I$ form a basis of \mathbb{R}^n . Two extreme points are *neighbors* if there exist bases for them, such that the intersection of the linear subspaces spanned by the two bases has dimension $n - 1$. The definition implies that also the line-segment between two neighboring extreme points is a face. Generically, this is a face of dimension 1, called *edge*. In the degenerate case, the two extreme points coincide, which happens if one extreme point is determined by two neighboring bases. In case P is of full dimension, the facets are of dimension $n - 1$, and each facet can be described by one inequality only. Therefore, if P is of full dimension, the system where the inequalities correspond one-to-one to the facets is the unique *minimal* defining system for P , up to multiplication of the inequalities by positive scalars.

Notice that the system of inequalities (*) that defines the class of convex games is of the type $Ax \geq 0$. Hence the class of convex games is a polyhedral cone in the vector space of games. Any strictly convex game is obviously in the interior of this cone, so this cone is of full dimension. (Existence of a strictly convex game was demonstrated by Shapley (1971), and this paper contains an example in Section 5.)

Also the core of a game v is obviously a polyhedron in \mathbb{R}^N , as the expression $x(S)$ for $S \subseteq N$ can be written as the linear expression $\langle e_S, x \rangle$, where e_S is the vector with ones at the coordinates of S and zeros elsewhere. We will abuse our terminology a bit, and say that a collection B of coalitions is a basis if $\{e_S \mid S \in B\}$ is a basis for \mathbb{R}^n .

4 Generalization of the Shapley-Ichiishi result

From Theorem 1 it follows that the set of limit games $L(v)$ for a strictly convex game v is the polyhedral cone of all convex games. In this section, we prove that the set $L(v)$ is a polyhedral cone for every balanced game v . In particular, we will describe how, given v , we can derive a system of linear inequalities whose solution set equals the polyhedral cone $L(v)$.

Let v be a balanced game. We say that a basis B of coalitions is *feasible for* v if there is an extreme point $x \in C(v)$ such that $B \subseteq T(v, x)$. We say that a coalition $T \subseteq N$ is a *neighbor* of B if there exists a feasible basis C for v such that $T \in C$ and $|B \cap C| = n - 1$. We say that T is *spanned by* B if the unique numbers λ_S for $S \in B$ to write

$$e_T = \sum_{S \in B} \lambda_S e_S$$

are nonnegative for all $S \in B \setminus \{N\}$.

To illustrate these concepts, consider the following example. Let v be the 3-player game defined by

$$v(S) = \begin{cases} 2 & \text{if } S = \{1\} \\ 2 & \text{if } S = \{2\} \\ 1 & \text{if } S = \{3\} \\ 2 & \text{if } S = \{1, 2\} \\ 5 & \text{if } S = \{1, 3\} \\ 4 & \text{if } S = \{2, 3\} \\ 10 & \text{if } S = N \end{cases}$$

There are 5 extreme points of $C(v)$. These are

$$\begin{aligned} a &= (2, 2, 6) \\ b &= (2, 5, 3) \\ c &= (4, 5, 1) \\ d &= (6, 3, 1) \\ e &= (6, 2, 2). \end{aligned}$$

For the 5 extreme core elements we see that

$$\begin{aligned} T(v, a) &= \{\{1\}, \{2\}, \{1, 2, 3\}\} \\ T(v, b) &= \{\{1\}, \{1, 3\}, \{1, 2, 3\}\} \\ T(v, c) &= \{\{3\}, \{1, 3\}, \{1, 2, 3\}\} \\ T(v, d) &= \{\{3\}, \{2, 3\}, \{1, 2, 3\}\} \\ T(v, e) &= \{\{2\}, \{2, 3\}, \{1, 2, 3\}\}. \end{aligned}$$

Now, for example, coalition $\{1, 3\}$ is a neighbor of $T(v, a)$, since $\{1, 3\} \in T(v, b)$, and a and b are neighboring extreme points. The coalition $\{1, 2\}$ is spanned by $T(v, a)$, since $e_{\{1,2\}} = e_{\{1\}} + e_{\{2\}}$. See also Figure 1.




Fig. 1. Core of the game v

Now, given the balanced game v , we construct a system of linear inequalities whose solution set equals the polyhedral cone $L(v)$. Let B be a feasible basis and let $T \subseteq N$ be a neighbor of B or spanned by B . Let $\lambda(S)$ for $S \in B$ be the unique real numbers such that

$$e_T = \sum_{S \in B} \lambda(S)e_S.$$

The linear inequality generated by the pair B, T is

$$w(T) \leq \sum_{S \in B} \lambda(S)w(S),$$

and we denote the set of all inequalities generated in this way by $I(v)$. Our claim is that $I(v)$ is a defining set for $L(v)$. In order to prove this claim we establish some preliminary facts.

Lemma 2 *Let v be a balanced game. If $T \in T(v, x) \setminus B$, where x is an extreme point of $C(v)$ and $B \subseteq T(v, x)$ is a basis, then the inequalities in $I(v)$ imply*

$$w(T) = \sum_{S \in B} \lambda(S)w(S).$$

Proof. Let x , B , and T be as above. Since T is a neighbor of B , $I(v)$ contains the inequality

$$w(T) \leq \sum_{S \in B} \lambda(S)w(S).$$

We show that the converse inequality is implied by the inequalities in $I(v)$. Consider the expression

$$e_T = \sum_{S \in B} \lambda(S)e_S.$$

There is at least one coalition $U \in B$ with $\lambda(U) > 0$. The collection $D := (B \setminus \{U\}) \cup \{T\}$ has exactly n elements and D is independent. Indeed, if D were not independent, then either B could not be a basis, or we could write e_T as a linear combination of the elements e_S with $S \in B$ without using e_U , which would violate the uniqueness of the coefficients. Consequently, D is a basis and U is a neighbor of D . Since

$$e_U = \frac{1}{\lambda(U)}e_T - \sum_{\substack{S \in B \\ S \neq U}} \frac{\lambda(S)}{\lambda(U)}e_S,$$

the inequality

$$w(U) \leq \frac{1}{\lambda(U)}w(T) - \sum_{\substack{S \in B \\ S \neq U}} \frac{\lambda(S)}{\lambda(U)}w(S)$$

is in $I(v)$. And since $\lambda(U) > 0$, this can be rewritten to

$$w(T) \geq \sum_{S \in B} \lambda(S)w(S),$$

which proves our claim. \square

For a game w that satisfies the inequalities in $I(v)$, we prove that w is a limit game for v . If x is an extreme point of $C(v)$ and B a basis for x , then the solution z to the system $y(S) = w(S)$ for all $S \in B$ is the obvious candidate for $z \in C(w)$ such that $T(v, x) \subseteq T(w, z)$. In the following lemma, we prove first that z indeed satisfies the requirement $T(v, x) \subseteq T(w, z)$, regardless whether $z \in C(w)$ or not.

Lemma 3 *Let v be a balanced game and let w be a game that satisfies the inequalities in $I(v)$. Let x be an extreme point in $C(v)$ and let B be a basis in $T(v, x)$. Let z be the unique solution to the system of equalities*

$$y(S) = w(S) \quad \text{for all } S \in B.$$

Then $T(v, x) \subseteq T(w, z)$.

Proof. Let $T \in T(v, x)$. We show that $T \in T(w, z)$. This is trivial if $T \in B$, so suppose that $T \notin B$. Write

$$e_T = \sum_{S \in B} \lambda(S) e_S.$$

By Lemma 2, the game w satisfies the equality

$$w(T) = \sum_{S \in B} \lambda(S) w(S),$$

hence

$$z(T) = \sum_{S \in B} \lambda(S) z(S) = \sum_{S \in B} \lambda(S) w(S) = w(T),$$

which proves our claim. \square

A geometrical interpretation of the following lemma is that, if w satisfies the system $I(v)$, the edges of $C(w)$ are parallel to those of $C(v)$, assuming that the candidate extreme points are indeed in $C(w)$.

Lemma 4 *Let v be a balanced game, and let w be a game that satisfies all inequalities in $I(v)$. Let x and \bar{x} be neighboring extreme points of $C(v)$, with feasible basis B and \bar{B} respectively, such that $|B \cap \bar{B}| = n - 1$. Let z, \bar{z} be defined respectively as the unique solution of*

$$y(S) = w(S) \quad \text{for all } S \in B$$

and

$$y(S) = w(S) \quad \text{for all } S \in \bar{B}.$$

Then there exists $\mu \geq 0$ such that

$$\bar{z} - z = \mu(\bar{x} - x).$$

Proof. First assume that $\bar{x} = x$. Then $\bar{z} = z$ by Lemma 3 and we can set $\mu = 0$. Now assume that $\bar{x} \neq x$. Write, for each $\alpha \in \mathbb{R}$,

$$z(\alpha) = z + \alpha(\bar{x} - x).$$

Let T be the unique coalition in $\bar{B} \setminus B$ and write

$$e_T = \sum_{S \in B} \lambda(S) e_S.$$

Since w satisfies all inequalities in $I(v)$ and T is a neighbor of B , in particular it satisfies the inequality

$$w(T) \leq \sum_{S \in B} \lambda(S) w(S).$$

Thus,

$$z(T) = \sum_{S \in B} \lambda(S) z(S) = \sum_{S \in B} \lambda(S) w(S) \geq w(T).$$

Furthermore, since $\bar{x} \neq x$, T is not tight at x and hence $\langle e_T, \bar{x} - x \rangle < 0$. This observation and the fact that $z(T) \geq w(T)$ implies that

$$\hat{\alpha} := \max\{\alpha \mid z(\alpha)(T) \geq w(T)\}$$

exists and is not negative. From the definition of $z(\alpha)$ it is now clear that we can set $\mu = \hat{\alpha}$ if we can show $z(\hat{\alpha}) = \bar{z}$.

To see why indeed $z(\hat{\alpha}) = \bar{z}$, notice that all coalitions $S \in B \cap \bar{B}$ are tight at all $z(\alpha)$ in the game w , because they are tight at z in w , and at both x and \bar{x} in v . Moreover, T is tight at $z(\hat{\alpha})$ in w by the definition of $\hat{\alpha}$. So, all coalitions in \bar{B} are tight at $z(\hat{\alpha})$ in w . Hence, $z(\hat{\alpha}) = \bar{z}$. \square

We have now developed enough equipment to prove the main theorem, namely that the system of inequalities in $I(v)$ precisely defines the set $L(v)$, which must therefore be a polyhedral cone.

Theorem 5 *Let v be a balanced game. Then a game w satisfies the inequalities in $I(v)$ if and only if $w \in L(v)$.*

Proof. A. Let $w \in L(v)$. We show that w satisfies the inequalities in $I(v)$. To this end, let B be a feasible basis for v , let $T \subseteq N$, and let $\lambda(S)$ for $S \in B$ be real numbers such that

$$e_T = \sum_{S \in B} \lambda(S) e_S.$$

Since w is a limit game for v , we can choose an extreme point $z \in C(w)$ such that $T(v, x) \subseteq T(w, z)$. Then

$$w(T) \leq z(T) = \langle z, e_T \rangle = \sum_{S \in B} \lambda(S) z(S) = \sum_{S \in B} \lambda(S) w(S).$$

This proves that w satisfies all inequalities in $I(v)$. Since we did not use the fact that T is a neighbor of B or spanned by B , in fact we proved that w satisfies many more inequalities.

B. Now suppose that w satisfies all inequalities in $I(v)$. Let x be an extreme point of $C(v)$. We need to prove that there exists a $z \in C(w)$ such that $T(v, x) \subseteq T(w, z)$. To this end, let $B \subseteq T(v, x)$ be a basis, and define z as the unique solution of the system

$$y(S) = w(S) \quad \text{for all } S \in B.$$

By Lemma 3, $T(v, x) \subseteq T(w, z)$, so it remains to prove that $z \in C(w)$.

First note that $z(N) = w(N)$, since $N \in T(v, x) \subseteq T(w, z)$. Now let $T \neq N$. We need to show that $z(T) \geq w(T)$. Consider the program

$$\mathcal{P} : \min \langle e_T, y \rangle \text{ s.t. } y \in C(v).$$

Solving this linear program by means of the simplex algorithm using starting point $y^1 = x$ and basis $B_1 = B$ for the initialization, yields a sequence

$$x = y^1, \dots, y^k =: \bar{x}$$

of extreme points of $C(v)$ together with a basis B_m for each point y^m . From the properties of the simplex algorithm we have

$$|B_{m+1} \cap B_m| = n - 1$$

and

$$\langle e_T, y^{m+1} - y^m \rangle \leq 0.$$

Now, for each B_m , let z^m be the unique solution to the system

$$y(S) = w(S) \quad \text{for all } S \in B_m$$

of linear equalities, and let $\bar{z} = z^k$. We will prove that $z(T) \geq \bar{z}(T) \geq w(T)$.

By Lemma 4, there exists nonnegative numbers μ_m , such that $z^{m+1} - z^m = \mu_m(y^{m+1} - y^m)$ for $m \in \{1, \dots, k\}$. Hence,

$$\begin{aligned} \bar{z}(T) = z^k(T) = \langle e_S, z^k \rangle &= \langle e_T, z^1 \rangle + \sum_{m=1}^{k-1} \langle e_T, z^{m+1} - z^m \rangle \\ &= z^1(T) + \sum_{m=1}^{k-1} \mu_m \langle e_T, y^{m+1} - y^m \rangle \\ &\leq z^1(T) = z(T). \end{aligned}$$

It remains to prove that $\bar{z}(T) \geq w(T)$. If $T \in T(v, \bar{x})$, then also $T(w, \bar{z})$, hence $\bar{z}(T) = w(T)$. Otherwise, consider the dual of the linear program \mathcal{P} , given by

$$\mathcal{D}: \quad \begin{array}{ll} \min & \sum_{S \neq \emptyset} \alpha_S v(S) \\ \text{subject to} & \sum_{S \neq \emptyset} \alpha_S e_S = e_T \\ & \alpha_S \geq 0 \quad \text{for all } S \neq \emptyset, N. \end{array}$$

Let λ_S ($S \neq \emptyset$) be an extreme optimal solution of the dual, and let B be the collection of coalitions S for which $\lambda_S \neq 0$. By the complementary slackness relations of linear programming, we have $\bar{y}(S) = v(S)$ for all S with $\lambda_S \neq 0$, hence $B \subseteq T(v, \bar{x})$. Since λ was chosen extreme, B is independent and contains at most n coalitions. If B is not already a basis, then we extend it to a basis with coalitions from $T(v, \bar{x})$. Then

$$e_T = \sum_{S \in B} \lambda_S e_S$$

with $\lambda_S \geq 0$ for all $S \in B \setminus \{N\}$, since λ is feasible for \mathcal{D} . We see that T is spanned by B , and since w satisfies all inequalities in $I(v)$, it satisfies in particular the inequality

$$w(T) \leq \sum_{S \in B} \lambda_S w(S).$$

Then

$$w(T) \leq \sum_{S \in B} \lambda_S w(S) = \sum_{S \in B} \lambda_S \bar{z}(S) = \bar{z}(T).$$

□

In the definition of $I(v)$, we tried to economize on the number of inequalities as much as possible. In case the polyhedral cone $L(v)$ defined by $I(v)$ is of full dimension, one might hope that $I(v)$ is the unique minimal defining system for it. We conclude this section with an example to show that this is not true in general. Consider the following game with player set $N = \{1, 2, 3, 4\}$.

$$v(S) = \begin{cases} 0 & \text{if } S = \{1\} \text{ or } S = \{4\} \\ 2 & \text{if } S = \{2\} \text{ or } S = \{3\} \\ 5 & \text{if } S = \{1, 2\} \text{ or } S = \{1, 3\} \text{ or } S = \{2, 3\} \\ -1 & \text{if } S = \{1, 4\} \text{ or } S = \{2, 4\} \text{ or } S = \{3, 4\} \\ -1 & \text{if } |S| = 3 \text{ and } 4 \in S \\ 9 & \text{if } S = \{1, 2, 3\} \\ 11 & \text{if } S = N. \end{cases}$$

With some effort, one can verify that $C(v)$ has twelve extreme points. These are

$$\begin{aligned} a &= (3, 4, 2, 2) & g &= (0, 5, 6, 0) \\ b &= (4, 3, 2, 2) & h &= (0, 6, 5, 0) \\ c &= (4, 2, 3, 2) & i &= (3, 2, 6, 0) \\ d &= (3, 2, 4, 2) & j &= (3, 6, 2, 0) \\ e &= (1, 4, 4, 2) & k &= (6, 2, 3, 0) \\ f &= (0, 5, 5, 1) & l &= (6, 3, 2, 0). \end{aligned}$$

For the twelve extreme core elements above we can check that

$$\begin{aligned} T(v, a) &= \{\{3\}, \{1, 3\}, \{1, 2, 3\}, N\} & T(v, g) &= \{\{1\}, \{4\}, \{1, 2\}, N\} \\ T(v, b) &= \{\{3\}, \{2, 3\}, \{1, 2, 3\}, N\} & T(v, h) &= \{\{1\}, \{4\}, \{1, 3\}, N\} \\ T(v, c) &= \{\{2\}, \{2, 3\}, \{1, 2, 3\}, N\} & T(v, i) &= \{\{2\}, \{4\}, \{1, 2\}, N\} \\ T(v, d) &= \{\{2\}, \{1, 2\}, \{1, 2, 3\}, N\} & T(v, j) &= \{\{3\}, \{4\}, \{1, 3\}, N\} \\ T(v, e) &= \{\{1, 2\}, \{1, 3\}, \{1, 2, 3\}, N\} & T(v, k) &= \{\{2\}, \{4\}, \{2, 3\}, N\} \\ T(v, f) &= \{\{1\}, \{1, 2\}, \{1, 3\}, N\} & T(v, l) &= \{\{3\}, \{4\}, \{2, 3\}, N\}. \end{aligned}$$

Notice that every collection of tight coalitions is a basis. It is then straightforward to prove that, for small perturbations of v , the core structure remains the same. Hence, the cone $L(v)$ is of full dimension, and has a unique minimal defining system. The defining system $I(v)$ however has redundant inequalities in it. To see this, consider the graph of neighboring extreme points, restricted to the six points a, b, c, d, e and f .




Fig. 2. Neighboring extreme points of $\text{Core}(v)$.

Each edge in this graph induces an inequality. For example, the inequality ($f \rightarrow e$) is generated by writing $e_{\{1,2,3\}}$ as a linear combination of the characteristic vectors corresponding to the elements in $T(v, f)$:

$$e_{\{1,2,3\}} = e_{\{1,2\}} + e_{\{1,3\}} - e_{\{1\}}.$$

The inequalities thus induced by the edges in this graph are

$$\begin{array}{lll} w(\{1, 2, 3\}) & \leq w(\{1, 2\}) + w(\{1, 3\}) - w(\{1\}) & (f \rightarrow e) \\ w(\{3\}) & \leq w(\{1, 2, 3\}) - w(\{1, 2\}) & (e \rightarrow a) \\ w(\{2, 3\}) & \leq w(\{1, 2, 3\}) + w(\{3\}) - w(\{1, 3\}) & (a \rightarrow b) \\ w(\{2\}) & \leq w(\{2, 3\}) - w(\{3\}) & (b \rightarrow c) \\ w(\{1, 2\}) & \leq w(\{1, 2, 3\}) - w(\{2, 3\}) + w(\{2\}) & (c \rightarrow d) \\ w(\{1, 3\}) & \leq w(\{1, 2, 3\}) - w(\{2\}) & (d \rightarrow e). \end{array}$$

Notice that the inequality ($e \rightarrow a$) is implied by inequalities ($b \rightarrow c$) and ($c \rightarrow d$). Symmetrically, ($d \rightarrow e$) is implied by ($a \rightarrow b$) and ($b \rightarrow c$).

5 Application to convex games

In this section, we investigate the consequences of Theorem 5, when it is applied to a strictly convex game. One could use Shapley's (1971) characterization of the core structure of these games, as described after Theorem 1, and apply Theorem 5 directly. We do not wish to rely on Shapley's result however. We will specify a strictly convex game and go through the whole process. Shapley's result makes clear that the result will be independent of our specific choice.

Define v^* by

$$v^*(S) := s^2$$

for each coalition S , where $s = |S|$. Then it is straightforward to prove that

$$v(S \cup T) + v(S \cap T) = v(S) + v(T) + 2ab,$$

where $a = |S \setminus T|$ and $b = |T \setminus S|$. Since $2ab \geq 0$, it follows that v^* satisfies the system (*). Indeed, v^* is strictly convex, since equality in (*) only occurs if $a = 0$ or $b = 0$, i.e. if $S \subseteq T$ or $T \subseteq S$.

We now establish the core structure of v^* .

Lemma 6 *The convex hull of the marginal vectors of v^* equals $C(v^*)$. Moreover, $T(v^*, m^\sigma(v^*)) = \{S_k^\sigma \mid k \in N\}$ for each marginal vector $m^\sigma(v^*)$.*

Proof. Let $m^\sigma(v^*)$ be a marginal vector of v^* . Assume w.l.o.g. that σ is the identity. Thus, the payoff to player i equals $i^2 - (i-1)^2 = 2i - 1$, and for a (non-empty) coalition S with $|S| = s$ we get that

$$m^\sigma(v^*)(S) = \sum_{i \in S} (2i - 1) \geq \sum_{i=1}^s (2i - 1) = 2 \cdot \frac{1}{2}s(s+1) - s = s^2 = v^*(S).$$

Let x be an extreme point of $C(v^*)$. To see that $T(v^*, x) = \{S_k^\sigma \mid k \in N\}$ for a suitably chosen permutation σ , it suffices to show that for any $S, T \subseteq N$ that are both tight at x in v^* , either $S \subseteq T$ or $T \subseteq S$. Note that

$$v^*(S \cap T) + v^*(S \cup T) \leq x(S \cap T) + x(S \cup T) = x(S) + x(T) = v^*(S) + v^*(T).$$

So, by the convexity of v^* , $v^*(S \cap T) + v^*(S \cup T) = v^*(S) + v^*(T)$. It follows that $S \subseteq T$ or $T \subseteq S$ because v^* is strictly convex. \square

Theorem 7 *The system $I(v^*)$ is the system of inequalities*

$$v(S \cup \{i, j\}) - v(S \cup \{j\}) \geq v(S \cup \{i\}) - v(S), \quad (***)$$

for all $i, j \in N$ ($i \neq j$) and all $S \subseteq N \setminus \{i, j\}$.

Proof. Let S, i, j be such that $S \subseteq N \setminus \{i, j\}$ and $i \neq j$. Let τ be a permutation for which the elements of S come first, then i , then j , and then the elements of $N \setminus (S \cup \{i, j\})$. Let σ be the permutation for which the elements of S are ordered as in τ , then j , then i , and then the elements of

$N \setminus (S \cup \{i, j\})$ as in τ . Note that $S \cup \{j\}$ is a neighbor of $T(v^*, m^\tau(v^*))$ and that $S, S \cup \{j\}, S \cup \{i, j\} \in T(v^*, m^\tau(v^*))$. Thus, since

$$e_T = e_{S \cup \{j\}} = e_{S \cup \{i, j\}} - e_{S \cup \{i\}} + e_S$$

(where e_S denotes the zero vector when S is empty), the inequality in $I(v^*)$ generated by the choice of S, i, j is

$$w(S \cup \{j\}) \leq w(S \cup \{i, j\}) - w(S \cup \{i\}) + w(S),$$

which can be rewritten as

$$w(S \cup \{i, j\}) - w(S \cup \{j\}) \geq w(S \cup \{i\}) - w(S)$$

So indeed, all inequalities in (***) are generated.

To see that no other inequalities are generated, associated with permutations σ, τ , note that the requirement $|T(v^*, m^\sigma(v^*)) \cap T(v^*, m^\tau(v^*))| = n - 1$ implies that τ can be obtained from σ by a switch of two players i, j , with j directly after i in σ . Then let S be the set of players that come before i in σ , and it is evident that σ and τ generate the inequality corresponding to S, i, j . \square

We obtain the following variant of the Shapley-Ichiishi result as a corollary.

Corollary 8 *A game v satisfies the inequalities in (***) if and only if all marginal vectors are elements of $C(v)$.*

Proof. Suppose that v satisfies the inequalities in (***). Then v is a limit game for v^* , by Theorem 5. Hence, $m^\sigma(v)$ is an extreme point of $C(v)$ for every σ .

Suppose that $m^\sigma(v) \in C(v)$ for every σ . Then note that every $m^\sigma(v)$ is extreme in $C(v)$, so it follows that v is a limit game for v^* . By Theorem 5, v satisfies the inequalities in $I(v^*)$, and by Theorem 7, this is the system (***). \square

Note that the set of inequalities (***) is the set $I(v)$ for *any* strictly convex game v . Hence, it also follows immediately that a game is convex precisely when it is a limit game of any strictly convex game.

Corollary 8 characterizes the games for which all marginal vectors are in the core with less inequalities than it is usually done. So, as a side result we proved that the system (***) implies the systems (**) and (*).

System (***), which we derived in Section 5, is in fact the *unique minimal representation* for the polyhedral cone of convex games, up to multiplication of the inequalities by positive scalars. To prove that each inequality in (***) indeed describes a facet of this cone, we will demonstrate that for each inequality in $I(v)$, there exist a game that violates the inequality, but satisfies all other inequalities in $I(v)$. In the proof, we make use of an alternative characterization of convex games by means of unanimity games.

A convenient basis for the vector space of all games was introduced in Shapley (1953). Let T be a non-empty coalition in the player set N . The corresponding *unanimity game* u_T is defined as follows.

$$u_T(S) = \begin{cases} 1 & \text{if } T \subseteq S \\ 0 & \text{otherwise.} \end{cases}$$

Shapley (1953) proved that the collection of unanimity games forms a basis of the vector space of all games. In other words, given a game v , there exist unique coefficients $(\alpha_T)_{T \neq \emptyset}$ such that

$$v = \sum_{T \neq \emptyset} \alpha_T u_T.$$

Many classes of games, like airport games (Littlechild and Owen, 1973) and sequencing games (Curiel *et al.*, 1989), can be characterized through restrictions on these coefficients. Also the class of convex games can be characterized through those coefficients.

Theorem 9 *The following two conditions are equivalent:*

- (a) *The game $v = \sum_{T \neq \emptyset} \alpha_T u_T$ is convex.*
- (b) *For all triplets (S, i, j) with $i \neq j$ and $i, j \notin S$:*

$$\sum_{T \subseteq S} \alpha_{T \cup \{i, j\}} \geq 0,$$

where the numbers α_T are the coefficients of the game v with respect to the basis of unanimity games.

Proof. By Corollary 8 it suffices to prove the equivalence of (***) and (b). Write $v = \sum_{T \neq \emptyset} \alpha_T u_T$. Let $i, j \in N$ such that $i \neq j$, and let $S \subseteq N \setminus \{i, j\}$.

Then

$$\begin{aligned}
v(S \cup \{i, j\}) - v(S \cup \{j\}) &\geq v(S \cup \{i\}) - v(S) \\
&\Leftrightarrow \\
[v(S \cup \{i, j\}) - v(S \cup \{j\})] - [v(S \cup \{i\}) - v(S)] &\geq 0 \\
&\Leftrightarrow \\
\left[\sum_{T \subseteq S \cup \{i, j\}} \alpha_T - \sum_{T \subseteq S \cup \{j\}} \alpha_T \right] - \left[\sum_{T \subseteq S \cup \{i\}} \alpha_T - \sum_{T \subseteq S} \alpha_T \right] &\geq 0 \\
&\Leftrightarrow \\
\sum_{T \subseteq S \cup \{j\}} \alpha_{T \cup \{i\}} - \sum_{T \subseteq S} \alpha_{T \cup \{i\}} &\geq 0 \\
&\Leftrightarrow \\
\sum_{T \subseteq S} \alpha_{T \cup \{i, j\}} &\geq 0.
\end{aligned}$$

□

Lemma 10 *Let $i, j \in N$ with $i \neq j$ and let $S \subseteq N \setminus \{i, j\}$. Let $v = \sum_T \alpha_T u_T$ with*

$$\alpha_T = \begin{cases} -1 & \text{if } T = S \cup \{i, j\} \\ 1 & \text{if } T = S \cup \{i\} \text{ or } T = S \cup \{j\} \text{ or } T \not\subseteq S \cup \{i, j\} \\ 0 & \text{otherwise.} \end{cases}$$

Then v violates the inequality

$$v(S \cup \{i, j\}) - v(S \cup \{j\}) \geq v(S \cup \{i\}) - v(S)$$

*and satisfies all other inequalities in (**).*

Proof. Notice that $\sum_{T \subseteq S} \alpha_{T \cup \{i, j\}} = -1 < 0$, so by Theorem 9, the condition for (S, i, j) is indeed violated. To show that all other inequalities are satisfied, let $k, l \in N$, $k \neq l$ and $V \subseteq N \setminus \{k, l\}$, such that the triplet $(V, k, l) \neq (S, i, j)$. Consider the quantity

$$\sum_{T \subseteq V} \alpha_{T \cup \{k, l\}}.$$

If $\alpha_{S \cup \{i, j\}}$ does not appear in the sum, then it is a sum of only nonnegative terms, hence it is nonnegative. If $\alpha_{S \cup \{i, j\}}$ does appear in the summation, discern two cases.

(a) $V \neq S$ and $\{i, j\} = \{k, l\}$. The fact that $\alpha_{S \cup \{i, j\}}$ appears in the above sum implies that the set S is a subset of V . So, since $V \neq S$, we can choose $m \in V \setminus S$. Then $\alpha_{\{m\} \cup \{k, l\}} = 1$ appears in the sum too, compensating for $\alpha_{S \cup \{i, j\}} = -1$ and consequently yielding a nonnegative outcome.

(b) $\{i, j\} \neq \{k, l\}$. Assume without loss of generality that $i \notin \{k, l\}$. Since $\alpha_{S \cup \{i, j\}} = -1$ appears in the sum, also $\alpha_{S \cup \{j\}} = 1$ appears in it, compensating the negative number and hence yielding a nonnegative outcome. \square

Corollary 11 *The system (***) is a facet description for the class of convex games.*

Proof. Since $L(v^*)$ is of full dimension, i.e., of dimension $2^n - 1$, we can choose a full-dimensional ball B of strictly convex games. Let v be a game that violates the inequality associated with (S, i, j) and none of the other inequalities in (**). Let K be the convex hull of v and the ball B . Let L be the intersection of K and the hyperplane $w(S \cup \{i, j\}) - w(S \cup \{j\}) = w(S \cup \{i\}) - w(S)$. It is obvious that L consists of convex games only, hence it is a subset of the face determined by (S, i, j) . It is also obvious that L has dimension $2^n - 2$, so it follows that (S, i, j) determines a face of dimension $2^n - 2$, hence a facet. \square

We conclude with a brief remark on the complexity of a minimal test for convexity. There are $\binom{n}{2}$ ways to choose two different players i and j in N , and 2^{n-2} ways to choose a coalition S that does not contain players i and j . Thus we have $2^{n-2} \binom{n}{2}$ inequalities of the form

$$v(S \cup \{i, j\}) - v(S \cup \{j\}) \geq v(S \cup \{i\}) - v(S).$$

Since the input size of a game is $x = 2^n - 1$ (i.e. the number of nonempty coalitions), the complexity of testing all inequalities in (***) is $\mathcal{O}(x(\log_2 x)^2)$. Testing the original system (*) has complexity $\mathcal{O}(x^2)$, and testing the system (**) of Ichiishi has complexity $\mathcal{O}(x^{\log_2 3} \log_2 x)$.

References

- Aumann R.J. and Maschler M., (1985) *Game theoretic analysis of a bankruptcy problem from the Talmud*, Journal of Economic Theory **36**, 195 – 213.
- Curiel I., Pederzoli G., and Tijs S., (1989) *Sequencing games*, European Journal of Operational Research **40**, 344 – 351.
- Driessen T.S.H., (1986), Contributions to the theory of cooperative games: the τ -value and k -convex games, *Ph.D Thesis, University of Nijmegen*.

- Granot D., Kuipers J., and Chopra S., (2002) *Cost allocation for a tree network with heterogeneous customers*, Mathematics of Operations Research **27**, 647 – 661.
- Granot D., Maschler M., Owen G., and Zhu W.R., (1996) *The kernel / nucleolus of a standard tree game*, International Journal of Game Theory **25**, 219 – 244.
- Hillas J. (1990) *On the definition of the strategic stability of equilibria*, Econometrica **58**, 1365 – 1390.
- Ichiishi T., (1981) *Supermodularity: applications to convex games and to the greedy algorithm for LP*, Journal of Economic Theory **25**, 283 – 286.
- Littlechild S.C. and Owen G., (1973) *A simple expression for the Shapley value in a special case*, Management Science **20**, 370 – 372.
- Megiddo N. (1978) *Computational complexity of the game theory approach to cost allocation for a tree*, Mathematics of Operations Research **3**, 189 – 196.
- O’Neill B., (1982) *A problem of rights arbitration from the Talmud*, Mathematical Social Sciences **2**, 345 – 371.
- Schrijver, A., (1986) Theory of linear and integer programming, *John Wiley and Sons*.
- Shapley L.S., (1953) *A value for n -person games*, in H. Kuhn and A.W. Tucker (eds.), Contributions to the Theory of Games II, Princeton University Press, 307 – 317.
- Shapley L.S., (1971) *Cores of convex games*, International Journal of Game Theory **1**, 11 – 26.
- Vermeulen A.J., Potters J.A.M., and Jansen M.J.M. (1996) *On quasi-stable sets*, International Journal of Game Theory **25**, 43 – 49.